

ATOMIC STRUCTURE
 $E = h\nu$
 $c = \lambda\nu$
 $\lambda = \frac{h}{mv}$
 $E_n = \frac{-2.178 \times 10^{-18}}{n^2}$ joule

EQUILIBRIUM

$K_a = \frac{[H^+][A^-]}{[HA]}$
 $K_b = \frac{[OH^-][HB^+]}{[B]}$
 $K_w = [OH^-][H^+] = 1.0 \times 10^{-14}$ @ 25°C
 $pH = -\log [H^+]$, $POH = -\log [OH^-]$
 $14 = pH + POH$
 $pH = pK_a + \log \frac{[A^-]}{[HA]}$
 $POH = pK_b + \log \frac{[B]}{[HB^+]}$
 $pK_a = -\log K_a$, $pK_b = -\log K_b$
 $K_p = K_c(RT)^{\Delta n}$
 where Δn = moles product gas - moles reactant gas

THERMOCHEMISTRY/KINETICS

$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$
 $\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$
 $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$
 $-RT \ln K = -2.303 RT \log Q$
 $q = m c \Delta T$
 $C_p = \frac{\Delta T}{\Delta H}$
 $\ln[A]_t - \ln[A]_0 = -kt$
 $\frac{1}{[A]_t} - \frac{1}{[A]_0} = kt$
 $\ln k = \frac{-E_a}{R} \left(\frac{1}{T} \right) + \ln A$

Faraday's constant, \mathcal{F} = 96,500 coulombs per mole of electrons
 Gas constant, $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$
 $= 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$
 $= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$

GASES, LIQUIDS, AND SOLUTIONS

$PV = nRT$
 $\left(P + \frac{n^2 a}{V^2} \right) (V - nb) = nRT$
 $P_{tot} = P_a + P_b + P_c + \dots$
 $P_a = P_{tot} \times X_a$, where $X_a = \frac{\text{moles } A}{\text{total moles}}$

$n = \frac{M}{m}$
 $K = C + 273$
 $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$
 $D = \frac{V}{m}$
 $n_{rms} = \sqrt{\frac{3RT}{M}}$
 $KE \text{ per molecule} = \frac{1}{2} m v^2$
 $KE \text{ per mole} = \frac{3}{2} RT$
 $\frac{1}{r_1} = \sqrt{\frac{M_2}{M_1}}$
 $\frac{1}{r_2} = \sqrt{\frac{M_1}{M_2}}$
 molality, M = moles solute per liter solution
 molality = moles solute per kilogram solvent
 $\Delta T_f = iK_f \times \text{molality}$
 $\Delta T_b = iK_b \times \text{molality}$
 $\pi = iMRT$
 $A = abc$

OXIDATION-REDUCTION; ELECTROCHEMISTRY

$Q = \frac{[C]^c [D]^d}{[A]^a [B]^b}$, where $aA + bB \rightarrow cC + dD$
 $I = \frac{I}{q}$
 $E_{cell} = E^\circ_{cell} - \frac{RT}{n\mathcal{F}} \ln Q = E^\circ_{cell} - \frac{0.0592}{n} \log Q$ @ 25°C
 $\log K = \frac{nE^\circ}{0.0592}$

P = pressure
 V = volume
 T = temperature
 n = number of moles
 D = density
 m = mass
 v = velocity

n_{rms} = root-mean-square speed
 KE = kinetic energy
 r = rate of effusion
 M = molar mass
 π = osmotic pressure
 i = van't Hoff factor
 K_f = molar freezing-point depression constant
 K_b = molar boiling-point elevation constant
 A = absorbance
 a = molar absorptivity
 b = path length
 c = concentration
 Q = reaction quotient
 I = current (amperes)
 q = charge (coulombs)
 t = time (seconds)
 E° = standard reduction potential
 K = equilibrium constant

Gas constant, $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$
 $= 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1}$
 $= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1}$
 Boltzmann's constant, $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$
 K_f for $H_2O = 1.86 \text{ K kg mol}^{-1}$
 K_b for $H_2O = 0.512 \text{ K kg mol}^{-1}$
 $1 \text{ atm} = 760 \text{ mm Hg}$
 $= 760 \text{ torr}$
 STP = 0.000°C and 1.000 atm
 Faraday's constant, $\mathcal{F} = 96,500$ coulombs per mole of electrons